

2005

Ecology of the tiger quoll *dasyurus maculatus maculatus* in coastal New South Wales

Deborah L. Andrew
University of Wollongong

Follow this and additional works at: <https://ro.uow.edu.au/theses>

University of Wollongong

Copyright Warning

You may print or download ONE copy of this document for the purpose of your own research or study. The University does not authorise you to copy, communicate or otherwise make available electronically to any other person any copyright material contained on this site.

You are reminded of the following: This work is copyright. Apart from any use permitted under the Copyright Act 1968, no part of this work may be reproduced by any process, nor may any other exclusive right be exercised, without the permission of the author. Copyright owners are entitled to take legal action against persons who infringe their copyright. A reproduction of material that is protected by copyright may be a copyright infringement. A court may impose penalties and award damages in relation to offences and infringements relating to copyright material.

Higher penalties may apply, and higher damages may be awarded, for offences and infringements involving the conversion of material into digital or electronic form.

Unless otherwise indicated, the views expressed in this thesis are those of the author and do not necessarily represent the views of the University of Wollongong.

Recommended Citation

Andrew, Deborah, Ecology of the tiger quoll *dasyurus maculatus maculatus* in coastal New South Wales, MSc thesis, School of Biological Sciences, University of Wollongong, 2005. <http://ro.uow.edu.au/theses/686>

NOTE

This online version of the thesis may have different page formatting and pagination from the paper copy held in the University of Wollongong Library.

UNIVERSITY OF WOLLONGONG

COPYRIGHT WARNING

You may print or download ONE copy of this document for the purpose of your own research or study. The University does not authorise you to copy, communicate or otherwise make available electronically to any other person any copyright material contained on this site. You are reminded of the following:

Copyright owners are entitled to take legal action against persons who infringe their copyright. A reproduction of material that is protected by copyright may be a copyright infringement. A court may impose penalties and award damages in relation to offences and infringements relating to copyright material. Higher penalties may apply, and higher damages may be awarded, for offences and infringements involving the conversion of material into digital or electronic form.

ECOLOGY OF THE TIGER QUOLL *DASYURUS MACULATUS*
MACULATUS IN COASTAL NEW SOUTH WALES

A thesis submitted in fulfillment of the requirements
for the award of the degree

MASTER OF SCIENCE (RESEARCH)

from

UNIVERSITY OF WOLLONGONG

by

DEBORAH L. ANDREW BSC. (HONS) UNSW

SCHOOL OF BIOLOGICAL SCIENCES

2005

CERTIFICATION

I, Deborah L. Andrew, declare that this thesis, submitted in fulfillment of the requirements for the award Master of Science (Research), in the School of Biological Sciences, University of Wollongong, is wholly my own work unless otherwise referenced or acknowledged. The document has not been submitted for qualifications at any other academic institution.

Deborah L. Andrew



Frontispiece: The Tiger Quoll *Dasyurus maculatus maculatus*

Abstract

The mammal fauna of Australia has undergone a dramatic and significant decline since the arrival and settlement of Europeans on the east coast of Australia. Eighteen native Australian mammals are considered extinct, 35 are endangered and 51 vulnerable, out of a total of 357 species. The Tiger Quoll is considered threatened in all mainland states, it is considered rare in Tasmania, and it is listed as endangered under Federal legislation. At the commencement of this study, little was known of the biology of the Tiger Quoll and virtually nothing was known of its ecology. In this study, I used captive Tiger Quolls at Featherdale Wildlife Park in Sydney, and wild populations at Limeburners Creek Nature Reserve on the mid-north coast of New South Wales and at Barren Grounds Nature Reserve and Budderoo National Park on the Illawarra escarpment south of Sydney to investigate several aspects of Tiger Quoll biology and ecology.

Studies on the breeding biology and growth of Tiger Quolls were undertaken at Featherdale Wildlife Park. Tiger Quolls have a single breeding season each year and at Featherdale Wildlife Park Tiger Quolls mated in May with births in June. Some females had a second and sometimes third oestrus cycle if the first didn't produce pouch young. Matings and births were fairly synchronous, occurring mostly over a two to four week period with a gestation length of about 21 days. Growth curves were developed for small pouch young and juveniles and age prediction curves were developed for small pouch young up to 60 days. Both males and females reached sexual maturity in 11 months and could breed in their first breeding season following their birth. Older females reared larger litters to teat release than one-year-old females. Females increased in weight prior to the mating period, developing fat deposits around the neck. Sexual dimorphism between the sexes, males being larger, became clearly apparent by 250 days of age.

In wild populations at Limeburners Creek Nature Reserve, mating occurred in June with the birth of a single litter recorded in July. Wild young were smaller than same-aged-captive young and growth of wild young is likely to be slower once they have left the pouch. Drought conditions are likely to affect the growth of young.

The spatial and social organisation of Tiger Quolls was investigated at Limeburners Creek Nature Reserve. Males and females were found to have large home range areas. Adult females had discrete home range areas, ranging in size from 509-1511 hectares, separate from other adult females, although some daughters occupied part of the mother's home range area. Adult males had much larger home ranges (2302-3401 hectares), which overlapped extensively with other males, and intersected 2-4 female home ranges. Den sharing was never observed, except between mothers and young, and sequential use of dens by different individuals only occurred several months after the former occupant had disappeared. Adult females generally moved to a new den each day except after pouch young released the teat and were deposited in a maternal den. Tiger Quolls undertook extensive movements on a daily and weekly basis, particularly males, with maximum recorded daily journeys of 5 and 8.5 kilometres for females and males respectively. Over longer periods males were recorded travelling over 20 kilometres in nine days. Activity was not exclusively nocturnal for adult females that may be active in early to mid-morning and late afternoon, especially in cooler months when nursing young.

Home ranges included a range of habitat types, heathland, sedgeland, shrubland, forest and woodland but the majority of locality records from trapping and radio-tracking were in forest and woodland. The majority of den sites were located within areas of eucalypt open forest and the majority of dens were eucalypt hollow logs or hollows in the trunks and limbs of live or dead standing trees. Dens were also located in burrows, which appeared to be preferred as maternal dens, under small buildings and in rocky headlands.

The diet of the Tiger Quoll at Limeburners Creek Nature Reserve and Barren Grounds Nature Reserve / Budderoo National Park (Illawarra) was predominantly carnivorous with a large proportion made up of vertebrate prey at both sites (85-91%). Mammals comprised the major vertebrate group with smaller contributions from birds (9-16%), reptiles (2-7%) and fish (13-24%). Insects were present in a greater proportion at Limeburners Creek Nature Reserve (26%) than the Illawarra area (13%). Plant material was present in the diet at both sites, as was garbage. There was potential for some competition between the Dingo and Tiger Quoll at Limeburners Creek Nature Reserve

as Dingoes took a large amount of bandicoot as prey, which was also a common prey item of Tiger Quolls.

Several aspects of the biology and ecology of the Tiger Quoll make it susceptible to a range of threats. Large home range size for both males and females makes the species vulnerable to clearing and fragmentation of forest and woodland habitats, to the threat of roadkill from increased traffic density and speed and to increased direct predation from eutherian carnivores such as the fox in rural landscapes and domestic pets in urban areas. Tiger Quolls are subject to direct persecution from humans due to their attraction to kill and eat caged birds such as chickens and their attraction to carrion increases the chances of death by roadkill.

The control of eutherian predators such as the fox and Dingo through 1080 poison baiting programs has the potential to affect Tiger Quoll populations through non-target kills and may have increased impact if undertaken in the restricted breeding season of Tiger Quolls and if undertaken for several consecutive years. Impacts may be greater on low-density populations. The removal of Dingoes may have a detrimental impact on Tiger Quoll populations if Dingoes are suppressing fox numbers. Further investigation of the potential impacts of poisoning programs and the relationships between eutherian and marsupial carnivores is required.

The protection of a number of large Tiger Quoll populations throughout the species geographic range and the protection and enhancement of forested corridors across the landscape may provide some buffer to potential impact of disease, habitat loss or degradation, or invasion of introduced species following climate change to help ensure the species conservation and possible recolonisation of areas of its former range.

Acknowledgements

I thank the Royal Zoological Society of New South Wales, Ethel Mary Read research fund for its contribution to the purchase of materials to construct hair sampling tubes. I thank Environment Australia (now Department of Environment and Heritage) for its grant to purchase cage traps and radio-tracking equipment.

I give my sincere thanks and gratitude to the staff and management of Featherdale Wildlife Park, especially Brad Walker, who allowed me to undertake trials of field methods on captive Tiger Quolls and studies of the growth and development of young Tiger Quolls. Brad's professional management of the captive Tiger Quoll population and his observations of Tiger Quolls, particularly over the breeding period, and assistance with measuring young Tiger Quolls made a very significant contribution to the breeding biology and growth chapter and his contribution is acknowledged throughout the document.

I thank the Manager and staff of the National Parks and Wildlife Service district office at Port Macquarie, especially Peter Clark, Cathy Mardell and Eric Claussen for their support and assistance during field studies at Limeburners Creek Nature Reserve and for the loan of a vehicle in the early stages of establishing the field project.

Liz Jeneid of Skyfarm Road, Jamberoo allowed me to collect Tiger Quoll scats from her property and Barbara Triggs of "Dead Finish", Genoa, Victoria provided expert identification of hair and bone remains in Tiger Quoll scats.

David Titley of Titley Electronics developed the first successful collar and radio transmitter package for wild Tiger Quolls for long-term monitoring, which allowed me to embark on the field study of home range, movements, den sites and social interactions of wild Tiger Quolls.

I thank my parents-in-law Hazel and Victor Hume for their enthusiastic support and encouragement and for feeding and housing me when I was undertaking field studies in Limeburners Creek Nature Reserve, and Victor, for his assistance and enthusiasm in

tracking down Tiger Quolls in the often thick and scrubby wilds of Limeburners Creek Nature Reserve. I wish to thank my late husband David Hume for his encouragement and support to embark on this endeavour to seek out new knowledge and his support with part of the field studies.

The residents of Point Plomer, especially Norm, assisted in monitoring Tiger Quolls that frequented that site, and oyster farmers Charlie Dick and Ray Bulley tolerated Tiger Quolls living under and around their huts and sheds and passed on valuable observations of Tiger Quoll behaviour.

I thank Michael Bedward of the NSW National Parks and Wildlife Service for his advice and assistance on several statistical matters. His assistance was greatly appreciated and kick-started the writing up process. The New South Wales National Parks and Wildlife Service provided access to computing resources and Sharon Davern, Peter Ewin and Kerry Oakes (NPWS) provided advice with compiling and formatting this document.

I thank my parents for their never-ending encouragement and support to complete this work and I thank my supervisor, Professor Rob Whelan for his support and encouragement to undertake the project and his ongoing tolerance and understanding to me in me completing this work over a difficult period in my life. I thank Rob for his enthusiastic encouragement to complete the work and advice and comments on drafts of this thesis.

Finally, but by no means least, I thank my friends and other members of my family for their support through this difficult period, without whose friendship, I would never have been able to complete this work.

CONTENTS

ABSTRACT	I
ACKNOWLEDGEMENTS	IV
1.0 INTRODUCTION.	1
1.1 Carnivorous marsupials–world distribution and diversity	1
1.2 Mammal decline in Australia	3
1.3 Distribution and conservation status of carnivorous marsupials of Australia and New Guinea	4
1.3.1 Tiger Quoll	4
1.3.2 Eastern Quoll	7
1.3.3 Western Quoll	9
1.3.4 Northern Quoll	9
1.3.5 New Guinea Quoll	10
1.3.6 Bronze Quoll	10
1.3.7 Tasmanian Devil	10
1.3.8 Thylacine	11
1.4 The Tiger Quoll–extent of knowledge of biology and ecology at commencement of this study	12
1.5 Morphology of Australasian carnivorous marsupials	12
1.6 Biology of Australasian carnivorous marsupials	14
1.6.1 Breeding	14
1.6.2 Longevity	16
1.7 Ecology of Australasian carnivorous marsupials	17
1.7.1 Diet	17
1.7.2 Social and spatial organisation	18
1.7.3 Dens and den sites	19
1.7.4 Habitat	20
1.8 Biology and ecology of the Tiger Quoll <i>Dasyurus maculatus</i> – gaps in knowledge	22
1.9 Aims of this study	23
2.0 STUDY AREAS.	25
2.1 Introduction	25
2.2 Featherdale Wildlife Park, Blacktown	25
2.3 Limeburners Creek Nature Reserve	28
2.4 Illawarra Escarpment, Jamberoo Pass	40
3.0 BREEDING BIOLOGY, GROWTH AND DEVELOPMENT OF THE TIGER QUOLL <i>DASYURUS MACULATUS MACULATUS</i>.	48

3.1	Introduction	48
3.2	Methods	50
3.2.1	Captive Tiger Quoll colony	50
3.2.2	Wild Tiger Quoll populations	53
3.3	Results	53
3.3.1	Captive Tiger Quoll colony	53
3.3.2	Wild Tiger Quoll populations	95
3.4	Discussion	117
3.4.1	Breeding season and reproductive cycle	117
3.4.2	Pouch development, marking and vocalisations	121
3.4.3	Litter size	122
3.4.4	Body measurements of wild Tiger Quolls	124
3.4.5	Weight changes in captive and wild Tiger Quolls	126
3.4.6	Growth of pouch young and juveniles	130
3.4.7	Morphological and behavioural development of young	134
4.0	SPATIAL AND SOCIAL ORGANISATION OF THE TIGER QUOLL <i>DASYURUS MACULATUS MACULATUS</i>.	138
4.1	Introduction	138
4.2	Methods	144
4.2.1	Study Area	144
4.2.2	Trapping and radio-tracking techniques	145
4.2.3	Home range analyses	152
4.2.4	Movement patterns	153
4.2.5	Proximity of individuals	153
4.2.6	Activity	153
4.3	Results	154
4.3.1	Home range estimates	154
4.3.2	Spatial and social organisation	163
4.3.3	Den use	185
4.3.4	Movement	191
4.3.5	Activity	198
4.4	Discussion	200
4.4.1	Home range area	200
4.4.2	Spatial and social organisation	205
4.4.3	Conclusion	221
5.0	DENS, DEN SITES AND HABITAT UTILISATION OF THE TIGER QUOLL <i>DASYURUS MACULATUS MACULATUS</i>.	223
5.1	Introduction	223
5.2	Methods	225
5.2.1	Dens	225
5.2.2	Habitat assessment	227
5.3	Results	229
5.3.1	Den types	229
5.3.2	Den spacing, distribution and site characteristics	251
5.3.3	Habitat utilisation	253

5.4	Discussion	258
5.4.1	Den types	258
5.4.2	Den use in other quoll species	260
5.4.3	Hollow formation in trees and logs	263
5.4.4	Function of dens	268
5.4.5	Habitat utilisation	271
5.4.6	Influence of other predators on habitat utilisation	274
6.0	DIET OF THE TIGER QUOLL <i>DASYURUS MACULATUS MACULATUS</i>.	278
6.1	Introduction	278
6.2	Methods	282
6.3	Results	288
6.3.1	Differences in the diet between study sites	288
6.3.2	Differences in the diet between years	296
6.3.3	Differences in the diet between seasons	296
6.3.4	Differences in the diet between males and females	297
6.3.5	Diet of the Dingo in Limeburners Creek Nature Reserve	309
6.4	Discussion	314
6.4.1	Comparison of the diet of the Tiger Quoll between study areas	314
6.4.2	Male and female diet	320
6.4.3	Diet of the Tiger Quoll throughout its range in eastern Australia	321
6.4.4	Potential competition between the Tiger Quoll and other predators	331
6.4.5	Comparison of the diet of the Tiger Quoll to other Australian quoll species and the Tasmanian Devil	335
7.0	SYNTHESIS - CONSERVATION AND MANAGEMENT OF THE TIGER QUOLL <i>DASYURUS MACULATUS MACULATUS</i> IN SOUTH-EASTERN AUSTRALIA.	339
7.1	Introduction	339
7.2	Biology and ecology of the Tiger Quoll- summary of findings of this study and other work	340
7.2.1	Breeding biology and growth of young	340
7.2.2	Longevity	342
7.2.3	Diet	342
7.2.4	Home range and spatial organisation	343
7.2.5	Movements and activity	344
7.2.6	Den requirements	345
7.2.7	Habitat	347
7.3	Potential threats to the conservation of the Tiger Quoll	348
7.3.1	Clearing and fragmentation of habitat	349
7.3.2	Degradation of habitat	353
7.3.3	Interactions with other mammalian predators	358
7.3.4	Potential impact of 1080 poisoning programs for the control of foxes, wild dogs and Dingoes and other introduced species on the conservation of the Tiger Quoll	362
7.3.5	Potential impact of trapping programs for the control of foxes, wild dogs and Dingoes	371
7.3.6	Potential impact of the expansion of the Cane Toad into south-eastern Australia	372
7.3.7	Disease and parasites	373
7.4	Conclusion	374

REFERENCES	376
APPENDICES	398

FIGURES

Figure 1.1: Distribution of Australian quolls of the genus <i>Dasyurus</i> .	5
Figure 1.2: Distribution of the Tiger Quoll in New South Wales.	6
Figure 2.1: Location of the main field study areas.	26
Figure 2.2: Climate graphs of the weather conditions recorded at North Parramatta.	27
Figure 2.3: Regional setting of Limeburners Creek Nature Reserve.	30
Figure 2.4: Climate graph of the weather conditions recorded at Port Macquarie.	31
Figure 2.5: Vegetation communities of Limeburners Creek Nature Reserve.	32
Figure 2.6: Regional setting of Barren Grounds Nature Reserve and Budderoo National Park.	41
Figure 2.7: Climate graphs for Moss Vale.	42
Figure 2.8: Vegetation communities of Barren Grounds Nature Reserve and Budderoo National Park.	44
Figure 3.1: Teat length of wild and captive female Tiger Quolls in relation to the age of the young.	62
Figure 3.2: Teat length of nursing wild and captive female Tiger Quolls in relation to month.	62
Figure 3.3: Crown-rump length of known-age litters regressed against age.	65
Figure 3.4: Head width of known-age litters regressed against age.	66
Figure 3.5: The regression of age predicted by crown-rump length for known-age litters.	67
Figure 3.6: The regression of age predicted by head width for known-age litters.	68
Figure 3.7: Head width of known-age and estimated-age litters to 345 days.	69
Figure 3.8: Head width of known-age and estimated-age litters to 556 days.	69
Figure 3.9: Crown-rump length of known-age and estimated-age litters to 168 days.	70
Figure 3.10: Nose-vent length of known-age and estimated-age litters to 281 days.	70
Figure 3.11: Tail length of known-age and estimated-age litters to 281 days.	71
Figure 3.12: Tail length of known-age and estimated-age litters to 556 days.	71
Figure 3.13: Tibia length of known-age and estimated-age litters to 281 days.	72
Figure 3.14: Tibia length of known-age and estimated-age litters to 556 days.	72
Figure 3.15: Weight of known-age and estimated-age litters to 780 days.	73
Figure 3.16: Weight changes in nursing females of litters 1, 2 and 3.	76
Figure 3.17: Weight changes in nursing females which were the mothers of Litters 4, 5, and 6.	77
Figure 3.18: Weight changes in nursing females which were the mothers of Litters 9, 10, 11 and 12.	78
Figure 3.19: Head width of two juvenile Tiger Quolls, <i>Banksia</i> (female) and <i>Bracken</i> (male).	106
Figure 3.20: Tail length of two juvenile Tiger Quolls, <i>Banksia</i> (female) and <i>Bracken</i> (male).	106
Figure 3.21: Tibia length of two juvenile Tiger Quolls, <i>Banksia</i> (female) and <i>Bracken</i> (male).	107
Figure 3.22: Weight of two juvenile Tiger Quolls, <i>Banksia</i> (female) and <i>Bracken</i> (male).	107
Figure 3.23: Monthly rainfall readings for the years 1990 to 1995 compared to average monthly rainfall for the Port Macquarie area.	111
Figure 3.24: Weight changes in adult and juvenile male and female Tiger Quolls trapped in the Limeburners Creek Nature Reserve.	113
Figure 4.1: The difference in the estimated position of the radio-collared Tiger Quoll to the actual position of the quoll once located.	150
Figure 4.2: The relationship of the mean linear error between the estimated position of the radio-collared Tiger Quoll to the length of the longest axis of the bearing triangle.	151
Figure 4.3: The distribution of the length of the long axis of the bearing triangle used to estimate a Tiger Quoll's position.	151
Figure 4.4a: Incremental increase in area of the home ranges of three adult female Tiger Quolls with increasing number of locations obtained.	156
Figure 4.4b: Incremental increases in home range area of three adult male Tiger Quolls with increasing number of locations.	156
Figure 4.5a: Incremental increases in home range area of three adult female Tiger Quolls with increasing number of monthly field trips.	157
Figure 4.5b: Incremental increases in home range area for three adult male Tiger Quolls with increasing number of monthly field trips.	157
Figure 4.6: Female Tiger Quoll home ranges in Limeburners Creek Nature Reserve.	159
Figure 4.7: Male Tiger Quoll home ranges in Limeburners Creek Nature Reserve.	160
Figure 4.8: Male and female home ranges showing den sites.	162
Figure 4.9: Tiger Quoll spatial organisation - Period 1.	171
Figure 4.10: Tiger Quoll spatial organisation -Period 2.	172
Figure 4.11: Tiger Quoll spatial organisation - Period 3.	173
Figure 4.12: Tiger Quoll spatial organisation - Period 4.	174
Figure 4.13: Tiger Quoll spatial organisation - Period 5.	175

Figure 4.14: Tiger Quoll spatial organisation - Period 6.	176
Figure 4.15: Tiger Quoll spatial organisation - Period 7.	177
Figure 4.16: The proximity of adult Tiger Quolls to each other.	184
Figure 4.17: The time of year when adult Tiger Quolls are within 500 m or less of another individual.	184
Figure 4.18: The number of days Tiger Quolls spent in the same den.	187
Figure 4.19: The period of the year and length of consecutive den stays by (a) female and (b) male Tiger Quolls.	188
Figure 4.20: Repeated use of dens by Tiger Quolls.	189
Figure 4.21: Mean daily movement of three female and three male Tiger Quolls over the year.	192
Figure 4.22: Mean daily movement over the year for individual (a) female and (b) male Tiger Quolls.	193
Figure 4.23: Weekly movement patterns of large male Tiger Quolls.	195
Figure 4.24: Weekly movement patterns of female Tiger Quolls.	196
Figure 4.25: Captures of Tiger Quolls in Limeburners Creek Nature Reserve throughout the year in respect to (a) total captures and (b) captures per hundred trap nights.	197
Figure 4.26: Diurnal activity in seven two hour time periods for (a) three female Tiger Quolls and (b) three male Tiger Quolls monitored throughout the year.	199
Figure 4.27: The relationship between body weight in grams (g) and home range area in hectares (ha) for male and female medium-sized Dasyurid insectivores and carnivores.	204
Figure 5.1: Den sites of adult male and female Tiger Quolls.	230
Figure 5.2: Den sites, locations and home ranges of Tiger Quolls of Limeburners Creek Nature Reserve in respect to vegetation structure.	231
Figure 5.3: The proportion of use of different den types by female and male Tiger Quolls.	232
Figure 5.4: The number of new dens located on each field trip compared to the number of den visits recorded.	234
Figure 5.5: Den types used by five females in respect to seasons of the year.	236
Figure 5.6: Size class of hollow logs used as dens by Tiger Quolls.	240
Figure 5.7: Size class and tree species of den logs used by Tiger Quolls.	240
Figure 5.8: The entrance diameter of hollow pipes at the base of logs used as dens by Tiger Quolls.	242
Figure 5.9: The relationship of the diameter of the hollow pipe at the base of logs used as dens to the size of the log as measured as diameter at breast height.	243
Figure 5.10: Size class of live trees as measured by diameter at breast height (dbh) that provided dens for Tiger Quolls.	245
Figure 6.1: Comparison of the percentage occurrence of prey and food items in the diet of the Tiger Quoll found in scats from Limeburners Creek Nature Reserve and the Illawarra escarpment.	292
Figure 6.2: Comparison of the percentage frequency of prey items in respect to the total number of prey items recorded in the diet of Tiger Quolls from Limeburners Creek Nature Reserve and the Illawarra escarpment.	292
Figure 6.3: Comparison of the percentage biomass contribution of prey items to the diet of Tiger Quolls from Limeburners Creek Nature Reserve and the Illawarra escarpment.	294
Figure 6.4: Prey and food items in the diet of Tiger Quolls from Limeburners Creek Nature Reserve in respect to season.	302
Figure 6.5: Comparison of the estimated biomass contribution of prey and food items to the diet of Tiger Quolls from Limeburners Creek Nature Reserve over seasons.	302
Figure 6.6: Prey and food items in the diet of Tiger Quolls from the Illawarra escarpment in respect to season.	305
Figure 6.7: Comparison of the estimated biomass contribution of prey and food items to the diet of Tiger Quolls from the Illawarra escarpment over seasons.	305
Figure 6.8: A comparison of the food and prey items in the diet of male and female Tiger Quolls from Limeburners Creek Nature Reserve.	308
Figure 6.9: A comparison of the estimated biomass contribution of food and prey items in the diet of male and female Tiger Quolls from Limeburners Creek Nature Reserve.	308
Figure 6.10: A comparison of measures, percentage occurrence in scats, percentage frequency of all prey items and percentage estimated biomass, in expressing the relative importance of prey items in the diet of the Dingo from Limeburners Creek Nature Reserve.	313
Figure 6.11: A comparison of prey items in the diet of the Dingo and Tiger Quoll from Limeburners Creek Nature Reserve.	313
Figure 6.12: A comparison of the diet of the Tiger Quoll from six sites in Australia.	325
Figure 6.13: A comparison of the diet of the Tiger Quoll throughout eastern Australia.	325
Figure 6.14: A comparison of broad food groups in the diet of four Australian quoll species.	334

TABLES

Table 3.1: Size and sex ratio of captive litters in respect to adult female Tiger Quoll age and weight.	60
Table 3.2: A comparison of body measurements of captive young Tiger Quolls.	74
Table 3.3: Morphological development of young Tiger Quolls.	80
Table 3.4: The date when captive young were first observed free of the teat.	82
Table 3.5: The greatest recorded measurements of wild Tiger Quolls.	96
Table 3.6: Age estimation of two young from <i>Christmas Bell's</i> litter.	99
Table 3.7: Body measurements of <i>Charlene's</i> young, <i>Banksia</i> (female) and <i>Bracken</i> (male).	99
Table 3.8: Litter size in wild Tiger Quolls as indicated by the number of expanded teats.	104
Table 3.9: Comparison of body measurements and age estimation of juvenile wild Tiger Quolls.	110
Table 3.10: Body weights and tail length of Tiger Quolls from south-eastern Australia.	125
Table 3.11: Morphological and behavioural development of the Western Quoll <i>Dasyurus geoffroii</i> , Eastern Quoll <i>D. viverrinus</i> and Northern Quoll <i>D. hallucatus</i> .	136
Table 4.1: Home range area for Tiger Quolls at Limeburners Creek Nature Reserve.	155
Table 4.2: Greatest linear distance between points in an individual's home range.	164
Table 4.3: The number of instances of sequential captures.	182
Table 4.4: The number of instances of repeated use of dens by Tiger Quolls.	186
Table 4.5: Home range size of medium and large sized insectivorous and carnivorous Dasyurid marsupials.	201
Table 5.1: Physical characteristics of dens used by Tiger Quolls.	237
Table 5.2: Tree species which provide dens to Tiger Quolls.	239
Table 5.3: Topographic position of dens used by Tiger Quolls.	252
Table 5.4: Vegetation type, height and cover at Tiger Quoll dens.	254
Table 5.5: Habitat use by Tiger Quolls in Limeburners Creek Nature Reserve.	256
Table 5.6: Tiger Quoll habitat use within their home range.	257
Table 6.1: Food and prey items recorded in the diet of Tiger Quolls from Limeburners Creek Nature Reserve and the Illawarra escarpment study sites.	291
Table 6.2: Food and prey items of Tiger Quolls from Limeburners Creek Nature Reserve and the Illawarra escarpment expressed as percentage biomass contribution to the diet.	293
Table 6.3: Food and prey items recorded in the diet of Tiger Quolls from Limeburners Creek Nature Reserve for the years 1993 and 1994.	298
Table 6.4: Food and prey items recorded in the diet of Tiger Quolls from the Illawarra escarpment for the years 1992, 1993 and 1994.	299
Table 6.5: Food and prey items recorded in the diet of Tiger Quolls from Limeburners Creek Nature Reserve by season.	300
Table 6.6: Food and prey items in the diet of Tiger Quolls from Limeburners Creek Nature Reserve in respect to season and biomass.	301
Table 6.7: Food and prey items recorded in the diet of Tiger Quolls from the Illawarra escarpment by season.	303
Table 6.8: Food and prey items in the diet of Tiger Quolls from the Illawarra escarpment in respect to season and biomass.	304
Table 6.9: Food and prey items recorded in the scats of adult male, adult female and juvenile Tiger Quolls from Limeburners Creek Nature Reserve.	306
Table 6.10: Biomass contribution of food and prey items recorded in the scats of adult male, adult female and juvenile Tiger Quolls from Limeburners Creek Nature Reserve.	307
Table 6.11: A comparison of the prey and food items recorded in the diet of the Dingo and Tiger Quoll from Limeburners Creek Nature Reserve.	311
Table 6.12: The estimated biomass contribution of prey items to the diet of the Dingo in Limeburners Creek Nature Reserve.	312

PLATES

Plate 1.1: Three species of Australian quolls.	8
Plate 2.1: Looking north east across the Limeburners Creek estuary.	35
Plate 2.2: Looking south from Big Hill headland to Point Plomer and Saltwater Lake.	35
Plate 2.3: Open forest of Blackbutt <i>Eucalyptus pilularis</i> and Pink Bloodwood <i>E. intermedia</i> .	36
Plate 2.4: Shrubland / heathland interface in the middle section of the Nature Reserve.	36
Plate 2.5: Heathland / sedgeland in the foreground with <i>Banksia</i> sp. and <i>Hakea</i> sp. with open forest of Scribbly Gum <i>Eucalyptus signata</i> and Needlebark Stringybark <i>E. planchoniana</i> on the low hill in the background.	37
Plate 2.6: Sedgeland in the foreground with taller <i>Gahnia</i> sp. prominent and tall swamp forest of Broad-leaved Paperbark <i>Melaleuca quinquenervia</i> on the skyline.	37
Plate 2.7: Eucalypt species that form canopy dominants in open forests of Limeburners Creek Nature Reserve.	38
Plate 2.8: Wetlands of Limeburners Creek Nature Reserve.	39
Plate 2.9: Looking south from Skyfarm Road along the Illawarra escarpment toward Barren Grounds Nature Reserve on the plateau.	46
Plate 2.10: Moist tall eucalypt forest and rainforest on the Illawarra escarpment near Macquarie Pass National Park.	46
Plate 2.11: Vegetation communities of the Illawarra escarpment and plateau.	47
Plate 3.1: Two year old female Tiger Quoll with thickened neck entering the breeding season.	55
Plate 3.2: Tiger Quolls mating.	55
Plate 3.3: Adult female Tiger Quoll during the mating season in May 1992 showing the expanded and reddened cloaca, distended pouch and bare patches on the abdomen.	57
Plate 3.4: Female Tiger Quoll in nest she has constructed.	61
Plate 3.5: Expanded teats in the pouch of a female Tiger Quoll.	61
Plate 3.6: Three newborn Tiger Quoll young of Litter 10.	83
Plate 3.7: Tiger Quoll young from Litter 4 aged 6 days old.	83
Plate 3.8: Tiger Quoll young from Litter 2 at 20 days old.	84
Plate 3.9: Tiger Quoll young from Litter 5 at 26 days old.	84
Plate 3.10: Tiger Quoll young from Litter 2 at 39 days old.	85
Plate 3.11: Tiger Quoll young from Litter 4 at 49 days old.	85
Plate 3.12: Tiger Quoll young of Litter 1 at 58 days old.	86
Plate 3.13: Tiger Quoll young of Litter 2 at 60 days old	86
Plate 3.14: Adult female Tiger Quoll with 60 day old young of Litter 2.	87
Plate 3.15: Tiger Quoll young of Litter 1 at 65 days old.	88
Plate 3.16: Tiger Quoll young of Litter 2 at 76 days old	88
Plate 3.17: Tiger Quoll young of Litter 1 at 84 days old.	89
Plate 3.18: Tiger Quoll young of Litter 1 at 84 days old climbing on their mothers back.	89
Plate 3.19: Tiger Quoll young of Litter 1 at 91 days old.	90
Plate 3.20: Tiger Quoll young of Litter 1 at 105 days old.	90
Plate 3.21: Tiger Quoll young of Litter 2 at 116 days old.	91
Plate 3.22: Tiger Quoll female young of Litter 1 at 147 days old.	91
Plate 3.23a: Female Tiger Quoll young from Litter 4 at 177 days old.	92
Plate 3.23b: Male Tiger Quoll young from Litter 4 at 177 days old.	92
Plate 3.24: Tiger Quoll male young from Litter 1 at 218 days old.	93
Plate 3.25: Adult female from Litter 1 at age 345 days old.	93
Plate 3.26: Adult female from Litter 6 in 1992 now 556 days old (~18months).	94
Plate 3.27: Adult female 914-920 days old (~29 months).	94
Plate 3.28a: Wild young female Tiger Quoll <i>Boronia</i> .	101
Plate 3.28b: Wild young male Tiger Quoll <i>Scribbly</i> .	101
Plate 3.29a: Young wild female Tiger Quoll <i>Banksia</i> .	102
Plate 3.29b: Young wild male Tiger Quoll <i>Bracken</i> .	102
Plate 4.1: Wild adult female Tiger Quolls of Limeburners Creek Nature Reserve.	169
Plate 4.2: Wild adult male Tiger Quolls of Limeburners Creek Nature Reserve.	170
Plate 5.1: Den logs used by Tiger Quolls.	247
Plate 5.2: Den under cabin used by male Tiger Quoll.	248
Plate 5.3: Log den used by female <i>Christmas Bell</i> when she was nursing young.	248
Plate 5.4: Tree dens used by Tiger Quolls in Limeburners Creek Nature Reserve.	249
Plate 5.5: Den burrow used by female Tiger Quoll <i>Raylene</i> .	250

Plate 5.6: Den burrow used by female Tiger Quoll <i>Charlene</i> .	250
Plate 6.1: Tiger Quoll scat deposition sites or latrine sites, on the Illawarra escarpment.	284
Plate 6.2: Scat deposition sites of captive Tiger Quolls at Featherdale Wildlife Park.	285